

OXALVAP

(Oxalic acid vaporizer)

INSTRUCTION MANUAL





Ing. Aleš Oharek Špitálská 738 684 01 Slavkov u Brna Czech Republic



www.oharkovi.cz



+420 776 181 982 +420 606 689 367



info@oharkovi.cz

What you need

- Acid vaporizer OXALVAP
- Power source (car battery 12 volts)
- Oxalic Acid crystals
- Measuring cup for crystals
- Electronic timer or stopwatch
- Eye protection protection glasses
- Mask approved for Oxalic Acid fumes (for example this one from 3M OV/N95)
- Acid resistant gloves
- Cloths to block entrances

Safety instructions

- Always wear your mask, eye protection and gloves
- Avoid skin contact with crystals, as this may cause burns
- More information on oxalic acid safety data sheet

Operating instructions

- 1. Add 1 gram per full sized box of Oxalic Acid crystals to the head of your Oxalic Acid Vaporizer. We use a 1/4 of a teaspoon, check your crystals with scales to ensure this is correct for you.
- 2. For example colony with two full sized boxes requires 2 grams for treatment.
- 3. Insert vaporizer into hive, either from the underneath the meshboard (if you can seal it) or from the front entrance.
- 4. Seal any ventilation with cloths.
- 5. Connect your vaporizer to the power source (12 volts) for two and a half minutes (maximum three minutes depends on battery capacity).
- 6. Disconnect power source.
- 7. Wait 3 to 5 minutes.
- 8. Removed all cloths and remove vaporizer.
- 9. Check that the vaporizer is cool before reuse or storage.

Conditions of use

- Carry out the treatment in still air at ambient temperatures above 5 °C.
- For better contact of bees with the substance, apply it in the evening.
- It can be applied only in the inter-bed period or in the case of compartments and swarms from which no honeycombs will be taken.
- The application does not work under the capped fetus.
- After 7 10 days, repeat the whole process at least 3 times.
- Repeat the procedure for three weeks, as acid fumes do not affect the mites in the fetal cells..
- The next day after treatment, count the number of mites on the hive pad. Mites fall down after application for another 2-3 weeks.
- Spring healing can be done only after the first flight of bees.

- Another application is recommended in the period from May to June.
- Autumn retreatment can only be performed during the period without bee brood in the months of November to December.
- The treatment should be combined with the evaporation of formic acid using equipment designed for this purpose.

NOTE: At lower ambient temperatures, for example in autumn, the fumes are more visible.

Technical parameters

- DIMENSIONS (LxHxW): 45cm x 1,2cm x 6cm
- WEIGHT: 250 g
- POWER SOURCE : car battery 12V
- MATERIAL : stainless steel, plastic

Package contents

- 1x vaporizer OXALVAP
- 1x handle
- 1x measuring cup for crystals
- 1x instruction manual

Notes